

Park City:
2200 Park Ave. Bldg D. Suite 100
Park City, Ut 84060

Old Mill Medical Center:
6360 So. 3000 East, Suite 210
Salt Lake City, UT 84121



HEIDEN ORTHOPEDICS

Eric Heiden, MD
Karen Heiden, MD
Corbett Winegar, MD
Jason Dickerson, DPM
Lindsey Marshall, PA-C
Phone: 435-615-8822

ERIC HEIDEN M.D.

ACL HAMSTRING AUTOGRAFT RECONSTRUCTION REHABILITATION PROTOCOL

	WEIGHT BEARING	BRACE	ROM	THERAPEUTIC EXERCISE
PHASE I 0-4 weeks	As tolerated with crutches*	0-2 weeks: locked in full extension for ambulation and sleeping	As tolerated	Heel slides, quad sets, patellar mobs, gastroc/soleus stretch**, SLR with brace in full extension until quad strength prevents extension lag***
PHASE II 4-6 weeks	Gradually discontinue crutch use	Discontinue use when patient has full extension and no extension lag	Maintain full extension and progressive flexion	Progress to weight bearing gastroc/soleus stretch and closed chain activities, begin hamstring stretching
PHASE III 6 weeks – 4 months	Full, without use of crutches and with a normalized gait pattern	None	Gain full and pain-free	Begin hamstring strengthening****, advance closed chain strengthening, progress proprioception activities, begin Stairmaster/Nordic Trac and running straight ahead at 12 weeks
PHASE IV 4-6 months	Full	None	Full and pain-free	Progress flexibility/strengthening, progression of function: forward/backward running, cutting, grapevine, etc., initiate plyometric program and sport-specific drills
PHASE V 6 months and beyond	Full	None	Full and pain-free	Gradual return to sports participation, maintenance program for strength and endurance At patient's discretion, a functional ACL brace may be used from 6 mo to 1 yr post-op

*Modified with concomitantly performed meniscus repair/transplantation or articular cartilage procedure

**This exercise is to be completed in a non-weight bearing position

***NO hamstring stretching until 4 weeks post-operative

****Avoid open chain quadriceps strengthening for first 4 months