

Park City:
2200 Park Ave. Bldg D. Suite 100
Park City, Ut 84060

Old Mill Medical Center:
6360 So. 3000 East, Suite 210
Salt Lake City, UT 84121



HEIDEN ORTHOPEDICS

Eric Heiden, MD
Karen Heiden, MD
Corbett Winegar, MD
Jason Dickerson, DPM
Lindsey Marshall, PA-C

Phone: 435-615-8822

ERIC HEIDEN M.D.

ARTHROSCOPIC SLAP REPAIR REHABILITATION PROTOCOL

	RANGE OF MOTION	IMMOBILIZER	THERAPEUTIC EXERCISE
PHASE I 0-4 weeks	Active/active-assisted: stretch to 40° of external rotation, 120° of forward flexion, internal rotation to stomach, abduction to 45° NO cross-body adduction	Worn at all times except for hygiene and therapeutic exercise	Elbow/wrist/hand ROM, grip strengthening, isometric in sling, external/internal rotation exercises with elbow at side No resisted forward flexion or biceps until 6 weeks post-op
PHASE II 4-8 weeks	Increase external rotation to 60° at side, forward flexion to 140°, internal rotation behind back to waist, abduction 60°	None	Advance isometrics in phase I to use of a theraband, continue with elbow/wrist/hand ROM and grip strengthening, begin prone extensions and scapular stabilizing exercises, gentle joint mobs Modalities prn
PHASE III 8-12 weeks	Progress to full active motion without discomfort Gentle passive stretching at end ranges	None	Advance theraband exercises to use of weights, continue with and progress exercises in phase II Include RTC, deltoid, and scapular stabilizers
PHASE IV 12 weeks – 6 months*	Full without discomfort	None	Advance exercises in phase III, begin functional progression to work/sport, return to previous activity level** Begin upper body ergometer, proprioception, plyometrics Return to throwing at 4 ½ months; throw from pitcher's mound at 6 months

*Patient may return to the weight room at 3 months, if appropriate

**Patient may return to competitive sports, including contact sports, by 6 months if approved